

A New Species of *Anagrus* (Hymenoptera: Mymaridae) from Tahiti, with Notes on Egg Parasitoids of Proconiine Sharpshooters (Hemiptera: Cicadellidae: Proconiini) in the World

Serguei V. Triapitsyn^{1,*}

Abstract

Triapitsyn, S. V. 2013. A new species of *Anagrus* (Hymenoptera: Mymaridae) from Tahiti, with notes on egg parasitoids of proconiine sharpshooters (Hemiptera: Cicadellidae: Proconiini) in the world. J. Taiwan Agric. Res. 62(4):289–299.

One new fairyfly (Hymenoptera: Mymaridae) species, *Anagrus* (*Anagrus*) *iti* sp. n., described from Tahiti Island, French Polynesia, was reared from eggs of the invasive glassy-winged sharpshooter, *Homalodisca vitripennis* (Germar) (Hemiptera: Cicadellidae), which is native to southeastern USA and northeastern Mexico. An overview of the taxonomic and biological studies on the egg parasitoids (Hymenoptera: Aphelinidae, Mymaridae, and Trichogrammatidae) of various proconiine sharpshooters (Cicadellidae: Cicadellinae: Proconiini) in the New World is given. Of particular importance are the natural enemies of *H. vitripennis*, a notorious vector of the phytopathogenic bacterium, *Xylella fastidiosa*; with reference to its egg parasitoids and biological control in the non-native range are also provided.

Key words: Leafhopper, Proconiini, Egg parasitoid, *Anagrus*, Biological control.

INTRODUCTION

Proconiine sharpshooters belong to the New World cicadelline leafhopper tribe Proconiini (Hemiptera: Cicadellidae: Cicadellinae). The most notorious species is the glassy-winged sharpshooter, *Homalodisca vitripennis* (Germar 1821), a vector of plant diseases caused by the phytopathogenic bacterium, *Xylella fastidiosa* (Blua *et al.* 1999). The glassy-winged sharpshooter is a self-introduced pest in California from the southeastern USA where it is native; it is also native to northeastern Mexico (Triapitsyn & Phillips 2000). Turner & Pollard (1959) provided a brief overview of

proconiine sharpshooter egg parasitoids (Hymenoptera: Mymaridae and Trichogrammatidae) in Georgia, USA. For a long time this was the only available publication on this group of natural enemies, which more recently has been shown to be amazingly diverse; almost every sharpshooter species has been found, upon closer examination, to have a complex of associated egg parasitoids. Establishment of *H. vitripennis* in California in the 1990s, later in the Hawaiian Islands and French Polynesia, and even more recently in Easter Island (Pilkington *et al.* 2005) prompted interest in proconiine sharpshooter investigations, including stud-

Received: July 30, 2013; Accepted: September 12, 2013.

* Corresponding author, e-mail: serguei.triapitsyn@ucr.edu

¹ Principal Museum Scientist, Department of Entomology, University of California, Riverside, CA, USA.

ies of their egg parasitoids in North America (Triapitsyn & Phillips 1996, 2000; Triapitsyn *et al.* 1998; Triapitsyn, Bezark *et al.* 2002; Triapitsyn, Hoddle *et al.* 2002), mainly for classical biological control purposes (Morgan *et al.* 2000, 2002; Triapitsyn & Hoddle 2001, 2002; Pilkington *et al.* 2004, 2005).

Grandgirard *et al.* (2007) conducted a thorough survey of the native egg parasitoids of the invasive *H. vitripennis* in French Polynesia; among them was an undescribed *Anagrus* sp. (Hymenoptera: Mymaridae) which was reared in Tahiti Island from an egg mass of the glassy-winged sharpshooter. It is described here as a new species, *A. iti*.

Due to the recent identification of *X. fastidiosa* in the vineyards of Taiwan (Su *et al.* 2013), which causes Pierce's disease of grapevines (the first such record in Asia), a summary of the known egg parasitoids in several families of Chalcidoidea (Hymenoptera) of the proconiine sharpshooter leafhoppers (its vectors) in the world is also given. Information on egg parasitoids of the leafhoppers from the tribe Cicadellini, also from the subfamily Cicadellinae, is scarce globally; these will be reviewed separately. Cicadellini contain several known vector species of *X. fastidiosa* in the USA and elsewhere; particularly, *Kolla paulula* (Walker 1858) is a potential vector of Pierce's disease of grapevine and a common cicadelline species in Taiwan (Shih *et al.* 2009; Su *et al.* 2013).

TAXONOMY: DESCRIPTION OF A NEW SPECIES OF MYMARIDAE

Anagrus (*Anagrus*) *iti* Triapitsyn, sp. n.
Anagrus sp.: Grandgirard *et al.* 2007:
183 (an undescribed species)

Type material. Holotype female on slide,

deposited in the Entomology Research Museum, University of California, Riverside, California, USA (UCRC), labeled: (1) 'FRENCH POLYNESIA: Society Islands, Tahiti I.[sland], Tapahi Point, 11.i.2005, J. Grandgirard, reared from eggs of Glassy-winged Sharpshooter, *Homalodisca coagulata*, on hibiscus'; (2) 'Mounted at UCR/ERM by V. V. Berezovskiy 2005 in Canada balsam' (UCRC ENT 265298). Paratypes: 16 females and 4 males on points, and 1 female and 1 male on slides (UCRC), same data as the holotype.

Description. Female. Body length (dry-mounted paratypes) 0.46–0.66 mm. Body pale to yellowish except frenum of scutellum white, basal gastral terga and exserted part of ovipositor sheaths brown; scape, pedicel, and legs yellowish, flagellum brownish; wings hyaline. Head about as wide as mesosoma. Antenna (Fig. 1A) with scape 2.9–3.5 times as long as wide, with cross-ridges, 2.1 times length of pedicel; F1 cylindrical, about half of pedicel length; F2 longest of funicle segments; F3 slightly shorter, F4–F6 subequal in length; longitudinal sensilla on F3 (1), F4 (1); F5 (1 or 2), and F6 (2); clava with 5 longitudinal sensilla, slightly shorter than combined length of two preceding segments. Mesosoma shorter than metasoma (Fig. 1B). Midlobe of mesoscutum with a pair of adnotaular setae. Fore wing (Fig. 1C) 8.3–8.8 times as long as wide, longest marginal seta 1.9–2.5 times maximum wing width; distal macrochaeta 1.7–1.8 times length of proximal macrochaeta; disc, besides a row of setae along each margin, with a single complete median row of setae originating behind marginal vein and a few additional, scattered setae, mostly at apex. Hind wing 26–27 times as long as wide, longest marginal seta 6.2–6.6 times maximum wing width; disc mostly bare except for admarginal rows of setae. Ovipositor notably exserted beyond apex of gaster (by




Fig. 1. *Anagrus (Anagrus) iti* (holotype female). (A) Antenna; (B) Metasoma; and (C) Fore wing.

0.15–0.16 times own length). External plates of ovipositor each with 3 distal setae. Ovipositor about 3.0 times length of protibia.

Measurements (mm) of the holotype female (as length or length : width). Body 0.805; head 0.154; mesosoma 0.258; gaster 0.406; ovipositor 0.375. Antenna: scape 0.094; pedicel 0.045; F1 0.024; F2 0.066; F3 0.063; F4 0.058; F5 0.055; F6 0.056; clava 0.103. Fore wing 0.581 : 0.07; longest marginal seta 0.133.

Hind wing 0.56 : 0.021; longest marginal seta 0.13.

Male. Body length (dry-mounted paratypes) 0.43–0.56 mm. Similar to female except for the normal sexually dimorphic features such as antenna and genitalia (Fig. 2A), and the following. Body notably darker than in female, with head and most of mesonotum light brown and gaster almost entirely brown to dark brown; fore wing (Fig. 2B) about 7.5 times as




Fig. 2. *Anagrus (Anagrus) iti* (paratype male). (A) Genitalia and (B) Fore wing.

long as wide, relatively more setose than in female.

Diagnosis. The new species belongs to the *incarnatus* species group of the nominate subgenus of *Anagrus*, as defined by Chiappini *et al.* (1996). Morphologically, it is most similar to the Australasian species *A. antipodus* Triapitsyn, 2001 and also to *A. oahuensis* Triapitsyn and Beardsley, 2000 known from the Hawaiian Islands and Mexico (Triapitsyn & Beardsley 2000; Triapitsyn 2002a). *Anagrus iti* has a longitudinal sensillum on F3 of female antenna, which *A. antipodus* lacks (Triapitsyn 2001), and 3 distal setae on the external plate of the ovipositor whereas *A. oahuensis* has 2 such setae.

Etymology. The species name ‘iti’ means ‘small’ in Tahitian language, referring to the minute size of the parasitoid.

Host. The type series of this species (18 females and 5 males) was reportedly reared from one egg mass of *H. vitripennis* on *Scae-*

vola sp. (Goodeniaceae) containing 8 eggs (Grandgirard *et al.* 2007) [as *Anagrus* sp.], although hibiscus (*Hibiscus* sp.; Malvaceae) was indicated on the labels; it is uncertain which is the correct host plant, and they would not be confused with one another.

NOTES ON THE KNOWN EGG PARASITOIDS OF PROCONIINI IN THE WORLD

Aphelinidae

***Centrodora* Förster, 1878:** Species of this genus appear to be more polyphagous egg parasitoids of some Hemiptera and are very difficult to identify. The two species reared from eggs of Proconiini remain undetermined: one *Centrodora* sp. from Tahiti and Moorea Islands, French Polynesia is a parasitoid of the non-native *H. vitripennis* (Grandgirard *et al.* 2007); the other *Centrodora* sp. was reared in Argentina from eggs of *Tapajosa rubromargin-*

ata Signoret, 1855 and *Tretogonia notatifrons* Melichar, 1926 (unpublished data).

Mymaridae

***Acmopolynema* Ogloblin, 1946:** One species of this genus, *A. sema* Schauff, was reared in Florida and Georgia, USA, from eggs of *Homalodisca insolita* (Walker 1858) (Triapitsyn, Hoddle *et al.* 2002).

***Anagrus* Haliday, 1833:** Besides *A. iti* from Tahiti described as a new species here, three other species of this genus have been recorded as egg parasitoids of Proconiini in the New World: *A. epos* Girault, 1911 in Minnesota, USA, a gregarious parasitoid of *Cuerna fenestella* Hamilton, 1970 (Hoddle & Triapitsyn 2004; Triapitsyn 2006), which was mass-reared and released without much success in California against *H. vitripennis* (Krugner *et al.* 2009; CDFA 2013), where its host specificity and other biological traits were studied by Krugner *et al.* (2008); *A. stethynioides* Triapitsyn, 2002, an occasional and poorly known parasitoid of *H. vitripennis* in Texas, USA (Triapitsyn 2006); and *A. breviphragma* Soyka, 1956 from *Dechacona missionum* (Berg 1879) in Argentina (Luft Albarracin *et al.* 2009).

***Gonatocerus* Nees ab Esenbeck, 1834:** This large and common genus contains many species that are known as egg parasitoids of various Proconiini, including those used in the biological control program against *H. vitripennis* in California (Peña *et al.* 2010; CDFA 2013), as summarized by Triapitsyn (2006) for the Nearctic region and Triapitsyn *et al.* (2010) for the Neotropical region. Details on their diversity, taxonomy, host associations and other biological traits are therefore omitted for brevity: these are readily available and too numerous to fit in this communication. All are solitary egg parasitoids, producing one wasp per host egg, except for *G. fasciatus* Girault, 1911 which is a gregarious parasitoid (Triapit-

syn *et al.* 2003). Biology of several species of *Gonatocerus* was thoroughly studied (Irvin & Hoddle, 2004, 2005a, 2005b; Virla *et al.* 2005; Son *et al.* 2012). They do not appear to be too host specific but rather are usually able to attack eggs of at least several genera and species of Proconiini (Boyd & Hoddle 2007; Logarzo *et al.* 2012; Lytle, Morse *et al.* 2012). All of them belong to the subgenus *G.* (*Cosmocomoidea* Howard 1908), which also contains a large number of undescribed species from Central and South America which are likely egg parasitoids of Proconiini (Triapitsyn 2002b; Triapitsyn *et al.* 2010), thus suggesting apparent co-evolution. Triapitsyn (2012) reported the following species of *Gonatocerus* from eggs of *H. vitripennis* in California: *G. ashmeadi* Girault, 1915, *G. fasciatus* (intentionally introduced), *G. incomptus* Huber, 1988, *G. morgani* Triapitsyn, 2006, *G. morrilli* (Howard 1908) (intentionally introduced), *G. novifasciatus* Girault, 1911, *G. triguttatus* Girault, 1916 (intentionally introduced), and *G. walkerjonesi* Triapitsyn, 2006. The southern and southeastern USA strains of *G. ashmeadi* were released in California against *H. vitripennis* (Morgan *et al.* 2002; Pilkington *et al.* 2005). *Gonatocerus ashmeadi* was found self-introduced in Oahu Island, Hawaii (USA), where it provides a good control of *H. vitripennis*; it was intentionally and successfully introduced into French Polynesia for very effective biological control against *H. vitripennis* (Grandgirard *et al.* 2008a, 2008b). It was also found self-introduced in Easter Island, Chile (Triapitsyn 2006). Paradell *et al.* (2012) summarized the known egg parasitoids of Proconiini in Argentina, the majority of which are *Gonatocerus* species described or revised taxonomically by Triapitsyn *et al.* (2006, 2007, 2008, 2010); *T. rubromarginata* is the most thoroughly studied host of many of these parasitoids.

Palaeoneura Waterhouse, 1915: One undetermined species of this genus was reared in Tahiti Island from eggs of *H. vitripennis* (Grandgirard *et al.* 2007).

Polynema Haliday, 1833: An undescribed species of *P. (Doriclytus)* sp. was reared in California, USA from eggs of *H. vitripennis*; it is only an occasional parasitoid of this host (Triapitsyn 2012). One or two other *P. (Doriclytus)* spp. were reared in Argentina from sentinel eggs of *T. rubromarginata* (G. A. Logarzo and S. V. Triapitsyn, unpublished).

Trichogrammatidae

Members of several genera are gregarious egg parasitoids of various Proconiini, all in the New World (Triapitsyn 2003); trichogrammatids are generally relatively more common on grassy vegetation and in dry habitats.

Burksiella De Santis, 1957: *Burksiella spirita* (Girault 1918) is a common egg parasitoid of *H. vitripennis* and also of *Oncometopia orbona* (Fabricius 1798) in the southeastern USA (Triapitsyn 2003 [as *Zagella spirita* (Girault)], 2012); a related form, which may or may not belong to this species, was reared in Montana, USA, from eggs of *Cuerna sayi* Nielson, 1965 (Triapitsyn *et al.* 2011). One or two different, undetermined species *Burksiella* sp.(p) was/were reported from eggs of *Homalodisca liturata* Ball, 1901 in Mexico (Triapitsyn & Bernal 2009). *Burksiella platensis* (De Santis 1957) is known from eggs of *T. rubromarginata* in Argentina (Triapitsyn 2003).

Ittys Girault, 1911: One undescribed species of this genus was reported from eggs of *H. liturata* in Mexico (Triapitsyn & Bernal 2009).

Oligosita Walker, 1851: An *Oligosita* sp., possibly *O. americana* Girault, 1909, was reared in Georgia from eggs of *H. insolita* (Turner & Pollard 1959; Triapitsyn 2003). Undetermined *Oligosita* spp. were also reared from eggs of two species of Proconiini in Ar-

gentina (Paradell *et al.* 2012); at least one of these is, however, a member of *Pseudoligosita* Girault, 1913 (Triapitsyn & Bernal 2009).

Paracentrobia Howard, 1897: *Paracentrobia acuminata* (Ashmead 1888) was reared in Florida and Georgia from eggs of *Cuerna costalis* (Fabricius 1803), *H. insolita*, and *H. vitripennis* (Turner and Pollard 1959; Triapitsyn 2003 [as *Paracentrobia (Brachistella)* sp.]; Tipping *et al.* 2005). Also *P. americana* (Girault 1917) is a parasitoid of *H. insolita* in Florida (Tipping *et al.* 2005). In Argentina, *P. tapajosae* Viggiani, 2009 is known from eggs of *T. rubromarginata* when Virla *et al.* (2009) studied biology of the parasitoids. Undetermined *Paracentrobia* spp. were also reared from eggs of three other species of Proconiini in Argentina (Paradell *et al.* 2012).

Pseudoligosita: Two species of this genus were reared from eggs of Proconiini: *P. plebeia* (Perkins 1912) from *H. liturata* in Mexico [Triapitsyn & Bernal 2009 (as *Pseudoligosita* sp.); Triapitsyn 2012] and also from eggs of *T. rubromarginata* in Argentina (Triapitsyn & Bernal 2009); and an undetermined and apparently undescribed *Pseudoligosita* sp. from eggs of *H. vitripennis* in California (Triapitsyn 2012). Biological traits of *P. plebeia*, which was reared under quarantine conditions in California on eggs of *H. vitripennis*, were reported by Triapitsyn & Bernal (2009) and Lytle, Bernal *et al.* (2012).

Ufens Girault, 1911: *Ufens ceratus* Owen, 2005 and *U. principalis* Owen, 2005 are common egg parasitoids of *H. liturata* and *H. vitripennis* in Mexico and USA (Al-Wahaibi *et al.* 2005; Triapitsyn & Bernal 2009; Triapitsyn 2012), and *U. ceratus* is also known from *Oncometopia clarior* (Walker 1851) in Mexico [Triapitsyn, Bezark *et al.* 2002 (as *Ufens* sp.)]. *Ufens niger* (Ashmead 1888) was reported from eggs of *C. costalis* and *Homalo-*

disca sp. (Turner & Pollard 1959; Triapitsyn 2003).

Zagella Girault, 1918: The sharpshooter *T. rubromarginata* is a known host of *Z. delicata* De Santis, 1970, its biology was reported by Logarzo *et al.* (2004). Undetermined *Zagella* spp. were also reared from eggs of three other species of Proconiini in Argentina (Paradell *et al.* 2012).

ACKNOWLEDGMENTS

I thank Dr. Hsien-Tzung Shih (TARI, Taiwan) for his kind advice, Dr. Julie Grandgirard (Service du Développement Rural, Papeete, Tahiti, French Polynesia) and Dr. Guillermo A. Logarzo (FuEDEI, Hurlingham, Buenos Aires, Argentina) for collecting the material and making it available for study, Mr. Vladimir V. Berezovskiy (UCRC) for mounting the specimens, and Dr. Douglas Yanega (UCRC) for his review of an early draft.

REFERENCES

- Al-Wahaibi, A. K., A. K. Owen, and J. G. Morse. 2005. Description and behavioural biology of two *Ufens* species (Hymenoptera: Trichogrammatidae), egg parasitoids of *Homalodisca* species (Hemiptera: Cicadellidae) in southern California. *Bull. Entomol. Res.* 95:275–288.
- Blua, M. J., P. A. Phillips, and R. A. Redak. 1999. A new sharpshooter threatens both crops and ornamentals. *Calif. Agric.* 53:22–25.
- Boyd, E. A. and M. S. Hoddle. 2007. Host specificity testing of *Gonatocerus* spp. egg-parasitoids used in a classical biological control program against *Homalodisca vitripennis*: A retrospective analysis for non-target impacts in southern California. *Biol. Contr.* 43:56–70.
- CDFA. 2013. GWSS biological control agents. (http://www.cdffa.ca.gov/pdcp/GWSS_Biological_Control_Agents.html).
- Chiappini, E., S. V. Triapitsyn, and A. Donev. 1996. Key to the Holarctic species of *Anagrus* Haliday (Hymenoptera: Mymaridae) with a review of the Nearctic and Palearctic (other than European) species and descriptions of new taxa. *J. Nat. Hist.* 30:551–595.
- Grandgirard, J., M. S. Hoddle, J. N. Petit, G. K. Roderick, and N. Davies. 2008a. Classical biological control of the glassy-winged sharpshooter, *Homalodisca vitripennis*, by the egg parasitoid *Gonatocerus ashmeadi* in the Society, Marquesas, and Austral Archipelagos of French Polynesia. *Biol. Contr.* 48:155–163.
- Grandgirard, J., M. S. Hoddle, J. N. Petit, G. K. Roderick, and N. Davies. 2008b. Engineering an invasion: classical biological control of the glassy-winged sharpshooter, *Homalodisca vitripennis*, by the egg parasitoid *Gonatocerus ashmeadi* in Tahiti and Moorea, French Polynesia. *Biol. Invasions* 10:135–148.
- Grandgirard, J., M. S. Hoddle, S. V. Triapitsyn, J. N. Petit, G. K. Roderick, and N. Davies. 2007. First records of *Gonatocerus dolichocerus* Ashmead, *Palaeoneura* sp., *Anagrus* sp. (Hymenoptera: Mymaridae), and *Centrodora* sp. (Hymenoptera: Aphelinidae) in French Polynesia, with notes on egg parasitism of the glassy-winged sharpshooter, *Homalodisca vitripennis* (Germar) (Hemiptera: Cicadellidae). *Pan-Pacific Entomol.* 83:177–184.
- Hoddle, M. S. and S. V. Triapitsyn. 2004. Searching for and collecting egg parasitoids of Glassy-winged Sharpshooter in the central and eastern USA. p.342–344. *in*: Proceedings of the 2004 Pierce's Disease Research Symposium. December, 7–10, 2004. Coronado Island Marriott Resort, CA, US. California Department of Food and Agriculture, Copeland Printing. Sacramento.
- Irvin, N. A. and M. S. Hoddle. 2004. Oviposition preference of *Homalodisca coagulata* for two *Citrus limon* cultivars and influence of host plant on parasitism by *Gonatocerus ashmeadi* and *G. triguttatus* (Hymenoptera: Mymaridae). *Florida Entomol.* 87:504–510.
- Irvin, N. A. and M. S. Hoddle. 2005a. Determination of *Homalodisca coagulata* (Hemiptera: Cicadellidae) egg ages suitable for oviposition by *Gonatocerus ashmeadi*, *Gonatocerus triguttatus*, and *Gonatocerus fasciatus* (Hymenoptera: Mymaridae). *Biol. Contr.* 32:391–400.
- Irvin, N. A. and M. S. Hoddle. 2005b. The competitive ability of three mymarid egg parasitoids (*Gonatocerus* spp.) for glassy-winged sharpshooter (*Homalodisca coagulata*) eggs. *Biol. Contr.* 34:204–214.
- Krugner, R., M. W. Johnson, D. J. W. Morgan, and J. G. Morse. 2009. Production of *Anagrus epos* Girault (Hymenoptera: Mymaridae) on *Homalodisca vitripennis* (Germar) (Hemiptera: Cicadellidae) eggs. *Biol. Contr.* 51:122–129.
- Krugner, R., M. W. Johnson, R. L. Groves, and J. G.

- Morse. 2008. Host specificity of *Anagrus epos*: a potential biological control agent of *Homalodisca vitripennis*. *BioControl* 56:439–449.
- Logarzo, G. A., E. G. Virla, E. Luft Albarracin, S. V. Triapitsyn, W. A. Jones, J. H. de León, and J. A. Briano. 2012. Host range of *Gonatocerus* sp. near *tuberculifemur* 'Clade 1' in Argentina, an egg parasitoid newly associated to the glassy-winged sharpshooter, *Homalodisca vitripennis* (Hemiptera: Cicadellidae), and candidate for its biological control in California, USA. *BioControl* 57:37–48.
- Logarzo, G. A., E. G. Virla, S. V. Triapitsyn, and W. A. Jones. 2004. Biology of *Zagella delicata* (Hymenoptera: Trichogrammatidae), an egg parasitoid of the sharpshooter *Tapajosa rubromarginata* (Hemiptera: Clypeorrhyncha: Cicadellidae) in Argentina. *Florida Entomol.* 87:511–516.
- Luft Albarracin, E., S. V. Triapitsyn, and E. G. Virla. 2009. Annotated key to the genera of Mymaridae (Hymenoptera: Chalcidoidea) in Argentina. *Zootaxa* 2129:1–28.
- Lytle, J. M., J. S. Bernal, and J. G. Morse. 2012. Biology of *Pseudoligosita plebeia* (Hymenoptera: Trichogrammatidae), an egg parasitoid of *Homalodisca* spp. (Hemiptera: Cicadellidae) collected from northwestern Mexico as a potential biocontrol agent of *H. vitripennis* in California. *J. Econ. Entomol.* 105:1532–1539.
- Lytle, J., J. G. Morse, and S. V. Triapitsyn. 2012. Biology and host specificity of *Gonatocerus deleoni* (Hymenoptera: Mymaridae), a potential biocontrol agent of *Homalodisca vitripennis* (Hemiptera: Cicadellidae) in California, USA. *BioControl* 57:61–69.
- Morgan, D. J. W., G. S. Simmons, L. M. Higgins, and K. Shea. 2002. Glassy-winged sharpshooter biological control in California: building framework for active adaptive management. p.140–143. *in*: Proceedings of the 3rd California Conference on Biological Control. August 15–16, 2002. Davis, CA, US. University of California, Davis.
- Morgan, D. J. W., S. V. Triapitsyn, R. A. Redak, L. G. Bezark, and M. S. Hoddle. 2000. Biological control of the glassy-winged sharpshooter: Current status and future potential. p.167–171. *in*: Proceedings of the California Conference on Biological Control II. July 11–12, 2000. Riverside, CA, US. Historic Mission Inn, Riverside.
- Paradell, S. L., E. G. Virla, G. A. Logarzo, and G. Delapé. 2012. Proconiini sharpshooters of Argentina, with notes on its distribution, host plants, and natural enemies. *J. Insect Sci.* 12:1–17.
- Peña, J. E., J. A. Jacas, S. Triapitsyn, B. J. Ulmer, and R. E. Duncan. 2010. Chapter 13. Biological control with egg parasitoids other than *Trichogramma* – The citrus and grape cases. p.341–371. *in*: Egg parasitoids in agroecosystems with emphasis on *Trichogramma*. (Cônoli, F. L., J. R. P. Parra, and R. A. Zucchi, eds.) Springer. Dordrecht, Netherlands. 479 pp.
- Pilkington, L. J., N. A. Irvin, E. A. Boyd, M. S. Hoddle, S. V. Triapitsyn, B. G. Carey, W. A. Jones, and D. J. W. Morgan. 2005. Introduced parasitic wasps could control glassy-winged sharpshooter. *Calif. Agric.* 59:223–228.
- Pilkington, L. J., N. Irvin, E. A. Boyd, M. S. Hoddle, S. V. Triapitsyn, B. G. Carey, W. A. Jones, and D. J. W. Morgan. 2004. Biological control of glassy-winged sharpshooter in California. p.133–136. *in*: Proceedings of the California Conference on Biological Control IV. July 13–15, 2004. Berkeley, CA, US. UC Berkeley, Berkeley.
- Shih, H. T., C. C. Su, C. Y. Feng, C. C. Fanjiang, W. F. Hung, and L. Y. Hung. 2009. Studies on the morphology, ecology, and host range for *Kolla paulula* (Walker, 1858) (Hemiptera: Membracoidea: Cicadellidae: Cicadellinae). *Formosan Entomol.* 29:353. (in Chinese with English abstract)
- Son, Y., H. Nadel, S. Baek, M. W. Johnson, and D. J. W. Morgan. 2012. Estimation of developmental parameters for adult emergence of *Gonatocerus morgani*, a novel egg parasitoid of the glassy-winged sharpshooter, and development of a degree-day model. *Biol. Contr.* 60:233–240.
- Su, C. C., C. J. Chang, C. M. Chang, H. T. Shih, K. C. Tzeng, F. J. Jan, C. W. Kao, and W. L. Deng. 2013. Pierce's disease of grapevines in Taiwan: Isolation, cultivation and pathogenicity of *Xylella fastidiosa*. *J. Phytopathol.* 161:389–396.
- Tipping, C., S. V. Triapitsyn, and R. F. Mizell, III. 2005. A new host record for the egg parasitoid *Paracentrobia americana* (Girault) (Hymenoptera: Trichogrammatidae) of the proconiine sharpshooter *Homalodisca insolita* (Walker) (Hemiptera: Clypeorrhyncha: Cicadellidae). *Florida Entomol.* 88:217–218.
- Triapitsyn, S. V. 2001. Review of the Australasian species of *Anagrus* (Hymenoptera Mymaridae). *Belgian J. Entomol.* 3:267–289.
- Triapitsyn, S. V. 2002a. Descriptive notes on a new and other little known species of *Anagrus* Haliday, 1833 (Hymenoptera: Mymaridae) from the New World tropics and subtropics. *Entomotropica* 17:213–223.
- Triapitsyn, S. V. 2002b. Species-level taxonomy of Mymaridae (Hymenoptera): current status and implications for biological control of leafhoppers of economic importance. p.89–94. *in*: Parasitic Wasps: Evolution, Systematics, Biodiversity and Biological Control. International Symposium: Parasitic Hyme-

- noptera: Taxonomy and Biological Control, Kőszeg, Hungary. (Melika, G. and C. Thuróczy, eds.), Agrofinform Kiadó és Nyomda Kft. Budapest. 480 pp.
- Triapitsyn, S. V. 2003. Taxonomic notes on the genera and species of Trichogrammatidae (Hymenoptera) - egg parasitoids of the proconiine sharpshooters (Hemiptera: Clypeorrhyncha: Cicadellidae: Proconiini) in southeastern USA. *Trans. Amer. Entomol. Soc.* 129:245–265.
- Triapitsyn, S. V. 2006. A key to the Mymaridae (Hymenoptera) egg parasitoids of proconiine sharpshooters (Hemiptera: Cicadellidae) in the Nearctic region, with description of two new species of *Gonatocerus*. *Zootaxa* 1203:1–38.
- Triapitsyn, S. V. 2012. Vouchering specimens of egg parasitoids of the glassy-winged sharpshooter collected by the CDFA pierce's disease biological control program in California and Texas A&M in Texas. p.8–11. *in: Pierce's Disease Research Progress Reports, December 2012.* (Esser, T. and R. Randhawa, eds.) California Department of Food and Agriculture. Sacramento, CA, US. 255 pp.
- Triapitsyn, S. V. and J. S. Bernal. 2009. Egg parasitoids of proconiini (Hemiptera: Cicadellidae) in northwestern Mexico, with description of a new species of *Gonatocerus* (Hymenoptera: Mymaridae). *J. Insect Sci.* 9:1–9.
- Triapitsyn, S. V. and J. W. Beardsley. 2000. A review of the Hawaiian species of *Anagrus* (Hymenoptera: Mymaridae). *Proc. Hawaii. Entomol. Soc.* 34:23–48.
- Triapitsyn, S. V. and M. S. Hoddle. 2001. Search for and collect egg parasitoids of glassy-winged sharpshooter in southeastern USA and northeastern Mexico. p.133–134. *in: Proceedings of the Pierce's Disease Research Symposium, December 5–7, 2001.* Coronado Island Marriott Resort, San Diego, CA, US. California Department of Food and Agriculture, Copeland Printing, Sacramento.
- Triapitsyn, S. V. and M. S. Hoddle. 2002. Search for and collect egg parasitoids of glassy-winged sharpshooter in southeastern USA and northeastern Mexico. p.94–95. *in: Proceedings of the Pierce's Disease Research Symposium, December 15–18, 2002.* Coronado Island Marriott Resort, San Diego, CA, US. California Department of Food and Agriculture, Digital Logistix, Sacramento.
- Triapitsyn, S. V. and P. A. Phillips. 1996. Egg parasitoid of glassy-winged sharpshooter. *Citrograph* 81:10.
- Triapitsyn, S. V. and P. A. Phillips. 2000. First host record of *Gonatocerus triguttatus* (Hymenoptera: Mymaridae) from eggs of *Homalodisca coagulata* (Homoptera: Cicadellidae), with notes on the distribution of the host. *Florida Entomol.* 83:200–203.
- Triapitsyn, S. V., D. B. Vickerman, J. M. Heraty, and G. A. Logarzo. 2006. A new species of *Gonatocerus* (Hymenoptera: Mymaridae) parasitic on proconiine sharpshooters (Hemiptera: Cicadellidae) in the New World. *Zootaxa* 1158:55–67.
- Triapitsyn, S. V., D. J. W. Morgan, M. S. Hoddle, and V. V. Berezovskiy. 2003. Observations on the biology of *Gonatocerus fasciatus* Girault (Hymenoptera: Mymaridae), egg parasitoid of *Homalodisca coagulata* (Say) and *Oncometopia orbona* (Fabricius) (Hemiptera: Clypeorrhyncha: Cicadellidae). *Pan-Pacific Entomol.* 79:75–76.
- Triapitsyn, S. V., G. A. Logarzo, E. G. Virla, and J. H. de León. 2007. A new species of *Gonatocerus* (Hymenoptera: Mymaridae) from Argentina, an egg parasitoid of *Tapajosa rubromarginata* (Hemiptera: Cicadellidae). *Zootaxa* 1619:61–68.
- Triapitsyn, S. V., G. A. Logarzo, J. H. de León, and E. G. Virla. 2008. A new *Gonatocerus* (Hymenoptera: Mymaridae) from Argentina, with taxonomic notes and molecular data on the *G. tuberculifemur* species complex. *Zootaxa* 1949:1–29.
- Triapitsyn, S. V., J. H. de León, and R. A. Rakitov. 2011. Egg parasitoid of the sharpshooter leafhopper *Cuerna sayi* (Hemiptera: Cicadellidae) in Montana, USA, with notes on the *Burksiella spirita* complex (Hymenoptera: Trichogrammatidae). *Boletín MIP (Manejo Integrado de Plagas, IMYZA-INTA, Argentina)* 21. (<http://anterior.inta.gov.ar/f/?url=http://anterior.inta.gov.ar/imyza/info/bol/mip/10/bol20/mip20.htm>).
- Triapitsyn, S. V., J. T. Huber, G. A. Logarzo, V. V. Berezovskiy, and D. A. Aquino. 2010. Review of *Gonatocerus* (Hymenoptera: Mymaridae) in the Neotropical region, with description of eleven new species. *Zootaxa* 2456:1–243.
- Triapitsyn, S. V., L. G. Bezark, and D. J. W. Morgan. 2002. Redescription of *Gonatocerus atriclavus* Girault (Hymenoptera: Mymaridae), with notes on other egg parasitoids of sharpshooters (Homoptera: Cicadellidae: Proconiini) in northeastern Mexico. *Pan-Pacific Entomol.* 78:34–42.
- Triapitsyn, S. V., M. S. Hoddle, and D. J. W. Morgan. 2002. A new distribution and host record for *Gonatocerus triguttatus* in Florida, with notes on *Acmopolynema sema* (Hymenoptera: Mymaridae). *Florida Entomol.* 85:654–655.
- Triapitsyn, S. V., R. F. Mizell, III, J. L. Bossart, and C. E. Carlton. 1998. Egg parasitoids of *Homalodisca coagulata* (Homoptera: Cicadellidae). *Florida Entomol.* 81:241–243.

- Turner, W. F. and H. N. Pollard. 1959. Life histories and behavior of five insect vectors of phony peach disease. U.S. Dept. Agriculture Tech. Bull. 1188:1-28.
- Virla, E. G., E. Luft Albarracin, S. V. Triapitsyn, G. Viggiani, and G. A. Logarzo. 2009. Description and biological traits of a new species of *Paracentrobia* (Hymenoptera: Trichogrammatidae), an egg parasitoid of the sharpshooter *Tapajosa rubromarginata* (Hemiptera: Cicadellidae) in Argentina. Stud. Neotr. Fauna Environ. 44:47-53.
- Virla, E. G., G. A. Logarzo, W. A. Jones, and S. Triapitsyn. 2005. Biology of *Gonatocerus tuberculifemur* (Hymenoptera: Mymaridae), an egg parasitoid of the sharpshooter, *Tapajosa rubromarginata* (Hemiptera: Cicadellidae). Florida Entomol. 88:67-71.

法屬大溪地島之 *Anagrus* 屬一新種 (膜翅目：縷小蜂科) 與 全球之脊大葉蟬族 (半翅目：葉蟬科：脊大葉蟬族) 卵寄生蜂記述

Serguei V. Triapitsyn^{1,*}

摘要

Triapitsyn I, S. V. 2013. 法屬大溪地島之 *Anagrus* 屬一新種 (膜翅目：縷小蜂科) 與全球之脊大葉蟬族 (半翅目：葉蟬科：脊大葉蟬族) 卵寄生蜂記述。台灣農業研究 62(4):289–299。

褐透翅尖頭大葉蟬 [*Homalodisca vitripennis* (Germar)] 原產於美國東南至墨西哥東北區域，目前已成為法屬波里尼西亞大溪地島的入侵物種。本研究描述採自大溪地島本種葉蟬蟲卵之縷小蜂科一個新種 – 微縷小蜂 [*Anagrus (Anagrus) iti* sp. n.]，並針對新世界區的脊大葉蟬族之卵寄生蜂 (包括膜翅目的蚜小蜂科、縷小蜂科及赤眼卵蜂科)，提供分類學與生物學的研究回顧。這部份又以傳播植物病原細菌 *Xylella fastidiosa* 最具效力的媒介昆蟲 - 褐透翅尖頭大葉蟬為重點，包括此種葉蟬在非原產地的卵寄生蜂與生物防治之相關研究。

關鍵詞：葉蟬、Proconiini、卵寄生蜂、*Anagrus*、生物防治。

投稿日期：2013年7月30日；接受日期：2013年9月12日。

* 通訊作者：serguei.triapitsyn@ucr.edu

¹ 美國加州大學河濱分校昆蟲學系首席科學研究員。美國 加州。